## Sowing Guideline : Broccoli & Cauliflower North West

VARIETY	JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEP	ОСТ	NOV	DEC
BROCCOLI												
Ares												
Naxos												
Parthenon												
Triton*												
CAULIFLOWER												
Alston												
Incline												
Merton												
Twister*												



Suggested sowing time: Sowing time applies to the area of seedling nursery and farm location

## Factors causing physiological disorders in cauliflower & broccoli

- Wrong sowing time
- Cold temperatures, especially below 7°C
- Excessive fertilisation of seedlings
- Cold grown seedlings
- Oversized seedlings at transplant
- Difference in temperatures between seedling nursery and farm
- Growth checks caused by heat, drought, water logging and disease

e-mail: info.saf@sakata.eu

website: www.sakata.co.za

Diurnal temperature swings

<u>Disclaimer:</u> This information is based on our observations and/or information from other sources. As crop performance depends on the interaction between the genetic potential of the seed, its physiological characteristics, and the environment, including management, we give no warranty express or implied, for the performance of crops relative to the information given nor do we accept any liability for any loss, direct or consequential, that may arise from whatsoever cause. Please read the Sakata Seed Southern Africa (Pty) Ltd Conditions of Sale before ordering seed.

Resistance: is the ability of a plant varieties under similar environmental conditions and pest or pathogen pressure. Resistant varieties may exhibit some disease symptoms or damage under heavy pest or pathogen pressure (HR = High resistance).

\* Experimental: This variety does not appear on the current South African Variety list, but has been submitted for registration. Recent version: Kindly contact Sakata or Area Representative for the most recent version of this Technical Bulletin.















Tel: +27 11 548 2800

Fax: +27 11 548 2820