## Sowing Guideline : Beetroot Western Cape

| VARIETY                                | JAN | FEB | MAR | APR | MAY | JUN | JUL | AUG | SEP | OCT | NOV | DEC |
|--|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| F1 HYBRID                              |     |     |     |     |     |     |     |     |     |     |     |     |
| Falcon                                 |     |     |     |     |     |     |     |     |     |     |     |     |
| Kestrel                                |     |     |     |     |     |     |     |     |     |     |     |     |
| Merlin                                 |     |     |     |     |     |     |     |     |     |     |     |     |
| Red Ace                                |     |     |     |     |     |     |     |     |     |     |     |     |
| SPECIALITY/NOVELTY                     |     |     |     |     |     |     |     |     |     |     |     |     |
| Chioggia Guardsmark<br>(Candy striped) |     |     |     |     |     |     |     |     |     |     |     |     |
| Touchstone Gold (Yellow)               |     |     |     |     |     |     |     |     |     |     |     |     |
| OPEN POLLINATED                        |     |     |     |     |     |     |     |     |     |     |     |     |
| Crimson Globe                          |     |     |     |     |     |     |     |     |     |     |     |     |
| Detroit Dark Red                       |     |     |     |     |     |     |     |     |     |     |     |     |
| Early Wonder                           |     |     |     |     |     |     |     |     |     |     |     |     |
| Globe Dark Red                         |     |     |     |     |     |     |     |     |     |     |     |     |



Suggested sowing time – sowing time depends on the production field's location

## Factors contributing to physiological disorders in beetroot

- Incorrect sowing time
- Incorrect sowing depth
- Excessive fertilisation
- Irregular irrigation
- Cold temperatures, especially below 7°C
- Growth stresses caused by heat, drought, water logging and disease
- Diurnal temperature flactuations this might induce bolting

<u>Disclaimer:</u> This information is based on our observations and/or information from other sources. As crop performance depends on the interaction between the genetic potential of the seed, its physiological characteristics, and the environment, including management, we give no warranty express or implied, for the performance of crops relative to the information given nor do we accept any liability for any loss, direct or consequential, that may arise from whatsoever cause. Please read the Sakata Seed Southern Africa (Pty) Ltd Conditions of Sale before ordering seed.

Resistance: is the ability of a plant varieties under similar environmental conditions and development of a specified pest or pathogen pressure. Resistant varieties may exhibit some disease symptoms or damage under heavy pest or pathogen pressure (HR = High resistance).

\* Experimental: This variety does not appear on the current South African Variety list, but has been submitted for registration. Recent version: Kindly contact Sakata or Area Representative for the most recent version of this Technical Bulletin.















Sakata Seed Southern Africa (Pty) Ltd. Copyright: (not to be reproduced)

Tel: +27 11 548 2800 Fax: +27 11 548 2820

e-mail: info.saf@sakata.eu

website: www.sakata.co.za