



OUTSTANDING QUALITIES

- ◆ INTERMEDIATE PINK ROOT RESISTANCE
- ◆ WIDELY ADAPTED
- ◆ MEDIUM TO LATE MATURITY

Texas Grano 502 PRR is a medium to late maturing short day brown onion with intermediate Pink root (Pt) resistance. The bulbs are uniform and have a granex shape. The scales are yellow, delicate and smooth. **Texas Grano 502 PRR** is well suited for early sowings in areas with a mild incidence of Pink root. Bulbs produced from transplants are much rounder than direct seeded bulbs. **Texas Grano 502 PRR** is a very high yielder and the bulbs have a refined neck. **Texas Grano 502 PRR** is suitable for most soil types.

SPECIAL VARIETAL REQUIREMENTS

- **Texas Grano 502 PRR** is well suited for production from direct sowing and transplants
- Grows longer than normal short day varieties, therefore need to apply follow-up fertiliser later than normal to ensure size and curing
- Contact area representative for a sowing guide

CHARACTERISTIC*	TEXAS GRANO 502 PRR
KIND	Onion (<i>Allium cepa</i> L.)
TYPE	Short day brown onion
MATURITY	Medium to late
BULB SHAPE	Granex
BULB SIZE	Medium to large (influenced by plant population)
BULB UNIFORMITY	Good
NECK SIZE	Thin
FIRMNESS	Medium to firm
BULB COLOUR	Yellow
FLESH COLOUR	White
SCALE RETENTION	Delicate
FLAVOUR	Mild
LEAF GROWTH HABIT	Upright
LEAF COLOUR	Dark green
BOLTING REACTION	Slow (with due regard to planting date)
DISEASE REACTION (SCIENTIFIC)	Intermediate resistance: <i>Pyrenochaeta terrestris</i> (Pt)
AVERAGE SEED COUNT	250 - 350 seeds per gram
POPULATION GUIDE	Final stand of 750 000 - 950 000 plants per ha
SEED REQUIREMENT	Direct sowing: 3.5 - 4.0 kg seed per ha Transplants: 5.0 - 5.5 kg seed per ha
STORAGE	Good
MARKETS / END USE	Fresh market and processing
SPECIAL FEATURES	Intermediate pink root resistance, widely adapted, high yield

* Characteristics given are affected by production methods such as soil type, nutrition, planting population, planting date and climatic conditions. Please read disclaimer.

Disclaimer: This information is based on our observations and/or information from other sources. As crop performance depends on the interaction between the genetic potential of the seed, its physiological characteristics, and the environment, including management, we give no warranty express or implied, for the performance of crops relative to the information given nor do we accept any liability for any loss, direct or consequential, that may arise from whatsoever cause. Please read the Sakata Seed Southern Africa (Pty) Ltd Conditions of Sale before ordering seed. **Resistance:** is the ability of a plant variety to restrict the growth and development of a specified pest or pathogen and/or the damage they cause when compared to susceptible plant varieties under similar environmental conditions and pest or pathogen pressure. Resistant varieties may exhibit some disease symptoms or damage under heavy pest or pathogen pressure (HR = High resistance, IR = Intermediate resistance).

* **Experimental:** This variety does not appear on the current South African Variety list, but has been submitted for registration.

Recent version: Kindly contact Sakata or Area Representative for the most recent version of this Technical Bulletin.





GENERAL TIPS FOR ONION PRODUCTION

Day length

All short day onion varieties require a photoperiod (day length) of between 10 - 12 h for bulb initiation; for instance, if a variety is exposed to a shorter than required photoperiod, there will be a high percentage of non-bulbing plants with thick necks. If a variety is exposed to a longer than required photoperiod, even only for a few days, the young plants will bulb prematurely, leading to reduced bulb size and yield.

Temperature

Even though day length is the primary factor responsible for bulb initiation (to which degree an onion variety will bulb), temperature plays a crucial role in the rate of bulbing. Onions need soil temperatures of between 15 - 25°C to germinate successfully. Temperatures of between 18 - 22°C is adequate for vegetative growth, however, bulbing will be slower at lower temperatures. Fast and proper bulbing requires temperatures of between 25 - 28°C. Temperatures below 10°C are detrimental and cold damage may occur, especially if the nitrogen available in the soil is low. When temperatures are below 10°C during the bulbing stage of the growth cycle, growth will be retarded and the plants might bolt (extreme fluctuations in day-night temperatures induce bolting).

Plant population

Plant population influences bulb size and therefore the final yield. The consumer generally prefers a medium-sized bulb (50 - 70 mm). The optimum plant population depends on the sought after/preferred marketable size, climate, soil type and planting date. The optimum plant population is between 750 000 – 950 000 plants per hectare (final stand).

Direct sowing: It is suggested to sow approximately 900 000 to 1 000 000 seed per hectare / 3.5 - 4.0kg seed per hectare, depending on the seed size and seed quality/germination factor (germination percentage of seed).

Transplanting and sets: Depending on the seed quality it is suggested to sow approximately 5.0 to 5.5kg seed for 1 hectare seedlings.

Nutrition

Onions are sensitive to waterlogging and require well-drained soil of at least 20 cm deep. The optimal soil pH for growing onions is between 5.5 and 6.8. A low soil pH increases the occurrence of certain soil-borne pathogens such as *Sclerotium cepivorum* (White rot) and *Fusarium oxysporum* f. sp. *cepea* (Basal rot).

The ratio of nitrogen (N) to potassium (K) is very important. In onion cultivation, potassium and sulphur play an important role in the keeping ability (storage and shelf life), firmness and health of onions. It is important to keep the ratio of nitrogen to potassium at 1:1.5. Thick necks in onions are not only an indication of incorrect sowing time, excessive nitrogen (applications), but might be due to an imbalance between nitrogen and potassium. All the phosphorus (P) and most of the potassium should be broadcasted before planting. The nitrogen component of the fertiliser programme should contain 45 - 50% ammonium nitrate (N-NH₄), which should be applied in the beginning of the growth phase.

Only nitrate-nitrogen (N-NO₃) should be applied in the final growth phase to prevent “green shouldering” on onions and to maintain continued growth in the cooler season. On alkaline soils, ammonium sulphate nitrate (ASN) should rather be used. Onions have a relatively shallow root system; therefore, it is important to apply nitrogen on a regular basis before bulbing (swelling of bulb/scale leaves) starts.

Sulphur, boron, zinc, molybdenum, magnesium, and potassium should always be available in adequate quantities; however, onions are sensitive to high levels of chloride.

As soon as onions tops start to collapse or “fall over”, all fertiliser applications should be stopped.

Onion curing

Adequate curing (windrowing or pyramid heaps) may require 2 - 4 weeks, depending on the weather (wind, temperature, humidity etc.) conditions. The best skin develops at 24 - 32°C. Curing can also be done by controlled ventilation in a storage room/warehouse by blowing heated air through the bottom of the onion pile to the top, at 9 - 15 cubic metres per minute per ton.

Onions are considered cured when the neck is tight and the outer scales are dry and make a rustling sound when handled. This condition is reached when onions have lost 3 - 5 % of their weight. If not adequately/properly cured, onions are likely to decay in storage.

Disease reaction definitions:

Resistance: is the ability of a plant variety to restrict the growth and development of a specified pest or pathogen and/or the damage they cause when compared to susceptible plant varieties under similar environmental conditions and pest or pathogen pressure. Resistant varieties may exhibit some disease symptoms or damage under heavy pest or pathogen pressure. Two levels of resistance are defined:

High/standard resistance (HR): plant varieties that highly restrict the growth and development of the specified pest or pathogen under normal pest or pathogen pressure when compared to susceptible varieties. These plant varieties may, however, exhibit some symptoms or damage under heavy pest or pathogen pressure.

Moderate/intermediate resistance (IR): plant varieties that restrict the growth and development of the specified pest or pathogen, but may exhibit a greater range of symptoms or damage compared to resistant varieties.

Moderately/intermediately resistant plant varieties will still show less severe symptoms or damage than susceptible plant varieties when grown under similar environmental conditions and/or pest or pathogen pressure.

Disclaimer: This information is based on our observations and/or information from other sources. As crop performance depends on the interaction between the genetic potential of the seed, its physiological characteristics, and the environment, including management, we give no warranty express or implied, for the performance of crops relative to the information given nor do we accept any liability for any loss, direct or consequential, that may arise from whatsoever cause. Please read the Sakata Seed Southern Africa (Pty) Ltd Conditions of Sale before ordering seed. **Resistance:** is the ability of a plant variety to restrict the growth and development of a specified pest or pathogen and/or the damage they cause when compared to susceptible plant varieties under similar environmental conditions and pest or pathogen pressure. Resistant varieties may exhibit some disease symptoms or damage under heavy pest or pathogen pressure (HR = High resistance, IR = Intermediate resistance).

* **Experimental:** This variety does not appear on the current South African Variety list, but has been submitted for registration.

Recent version: Kindly contact Sakata or Area Representative for the most recent version of this Technical Bulletin.

